

GST-Tag Mouse Monoclonal Antibody(Q16)

Catalog	TDY030C	TDY030F	Tel: 010-82908854
			Free: 400-0620-621
Quantity	50μL	100μL	Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB	N/A	N/A	IgG1

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% **sodium azide** as Preservative and 50% Glycerol.
Store at **-20°C**. **Do not aliquot the antibody.**


Recommended dilutions: WB: 1:2,000-5,000

Optimal dilutions should be determined by the end user.

Specificity: The GST tag antibody can recognize C-terminal, internal, and N-terminal GST fusion proteins.

Alternative Names: GST antibody

Background: The glutathione S-transferase (GST, previously known as ligandins) family of enzymes are composed of many cytosolic, mitochondrial, and microsomal proteins. GSTs are present in eukaryotes and in prokaryotes, where they catalyze a variety of reactions and accept endogenous and xenobiotic substrates. GST tag antibody can be helpful in detecting the fusion protein during purification as well as the cleavage of GST from the protein of interest. GST tag antibody has wide applications that could include your research on GST proteins or GST fusion recombinant proteins.



Western blot analysis of 0.2ug GST protein with GST (Q16) Mouse mAb
diluted at 1:5,000