

天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

Bovine anti-Goat IgG (H+L), HRP Conjugated

Catalog S008F Tel: 010-82908854

Free: 400-0620-621

Quantity 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC	Goat	N/A	IgG

Storage Buffer & Condition: PBS, pH 7.4,15mg/ml Bovine Serum Albumin as Stablizer and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.

Recommended dilutions: WB: 1:2,000-5,000 IHC: 1:100-500

Optimal dilutions should be determined by the end user.

Specificity: The antibody reacts with whole molecule goat IgG. The antibody minimal cross-reaction with **bovine**, **chicken**, **guinea pig**, **Syrian hamster**, **horse**, **mouse**, **rabbit**, **and rat** serum proteins, but it may cross-react with immunoglobulins from other species.

Note: Avoid use of Sodium azide as a preservative, it will substantially inhibit the enzyme activity of *HRP*.

Applications: WB-Western blot IHC-Immunochemistry IF-Immunofluorescence IP-Immunoprecipitation ChIP-Chormatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig