

NF κ B p65 Mouse Monoclonal Antibody (14H2)

Catalog TDY076C TDY076F

Tel: 010-82908854

Quantity 50 μ L 100 μ L

Free: 400-0620-621

Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC, IP	H,M,R	~65KD	IgG1

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% **sodium azide** as Preservative and 50% Glycerol.


Store at **-20°C**. **Do not aliquot the antibody.**

Recommended dilutions: WB: 1:1,000-3,000 IHC: 1:200 IP:1:200


Optimal dilutions should be determined by the end user.

Specificity: NF κ B p65 Mouse monoclonal antibody detects endogenous p65 proteins.


Background: NF κ B p65 is ubiquitinated leading to its proteosomal degradation, which is required for termination of the NF κ B response. Phosphorylation of NF κ B p65 on S536 stimulates acetylation of K310 by CBP, enhancing transcriptional activity. NF κ B p65 is also acetylated at K122, enhancing DNA binding and impairing the interaction with NFKBIA. The protein is deacetylated by HDAC3. Invasion of a host by a pathogen is frequently associated with the activation of NF- κ B, which coordinates various aspects of immune function required for resistance to infection.




Western blot analysis of 1) Hela, 2)293T,3)HepG with NF κ B p65 mAb diluted at 1:2,000.



Western blot analysis of 1) Mouse Heart, 2)Rat Heart,3)Rat Liver, 4) Rat Spleen with NF κ B p65 mAb diluted at 1:3,000.



IHC staining of mouse hippocampus tissue with NF κ B p65 mouse mAb(14H2) diluted at 1:200.



1, Input: Hela Cell Lysate

2, IP product: IP dilute 1:200

Western blot analysis: primary antibody : TDY076 1:2,000

Secondary antibody: Goat anti-Mouse IgG, Light chain specific(S003), 1:5,000

Applications: WB-Western blot IHC-Immunocytochemistry IF-Immunofluorescence IP-Immunoprecipitation ChIP-Chromatin Immunoprecipitation
Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig