

天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

β -actin Mouse Monoclonal Antibody(A02)

Catalog TDY045C TDY045F Tel, 010-82908854

Free: 400-0620-621

Quantity 50µL 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB. IHC. IF	H, R, M, Mk, Dg, Ch, Hm, Rb, Pg, Sh	43KD	laG1

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:5,000-10,000 IHC: 1:200 IF: 1:200


Optimal dilutions should be determined by the end user.

Specificity: Antibody can detects endogenous β-actin protein.

Alternative Names: actin antibody, b-actin antibody, beta-actin antibody

Background: β -actin is one of six different actin isoforms that have been identified. The actin molecules found in cells of various species and tissues tend to be very similar in their immunological and physical properties. Therefore, antibodies against β -actin are useful as loading controls for Western Blotting. However it should be noted that levels of β -actin may not be stable in certain cells. For example, expression of β -actin in adipose tissue is very low and therefore β -actin should not be used as loading control for these tissues.





Western blot analysis of Hela (1), Rat brain (2), Mouse brain(3), with β -actin IF analysis of Hela with TDY045 diluted at 1:500. mouse mAb(A02) diluted at 1:10,000.