

天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

Histone H3 (mono methyl K79) Mouse Monoclonal Antibody(5G5)

Catalog TDY120C TDY120F Tel: 010-82908854

Free: 400-0620-621

Quantity 50µL 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC	H, R, M	15KD	lgG1

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.


Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:500-2,000 IHC: 1:200-500

Optimal dilutions should be determined by the end user.

Specificity: Antibody can detects endogenous Histone H3 (mono methyl K79) protein. **Alternative Names:** H3 histone antibody, HIST1H3A antibody, Histone cluster 1, H3a antibody

Background: Histone H3 is one of the five main histone proteins involved in the structure of chromatin in eukaryotic cells. Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.






Western blot analysis of Hela with Histone H3 (mono methyl K79) (5G5) Mouse MAb diluted at Western blot analysis of 1) Rat Testis tissue, 2) Raw264.7 with Histone H3 (mono methyl K79) (5G5) Mouse MAb diluted at 1:2000.

1) 1:1,000

2) 1:3,000



Immunohistochemical analysis of paraffin-embedded Human Skin using Histone H3(momo methyl K79) (TDY120) Mouse mAb diluted at 1:500.

Applications: WB-Western blot IHC-Immunochemistry IF-Immunofluorescence IP-Immunoprecipitation ChIP-Chormatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig