天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

Histone H3(tri methyl K9) Mouse Monoclonal Antibody(4H8)

Catalog TDY136C TDY136F Tel, 010-82908854

Free: 400-0620-621

Quantity 50µL 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IP, IHC	H, R, M	15KD	lgG1

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:1,000-3,000 IP:1:200 IHC: 1:200-500

Optimal dilutions should be determined by the end user.

Specificity: The Histone H3(tri methyl K9) antibody can detects endogenous Histone H3(tri methyl K9) protein.

Alternative Names: H3 histone antibody, HIST1H3A antibody, Histone cluster 1, H3a antibody


Background: Histone H3 is one of the five main histone proteins involved in the structure of chromatin in eukaryotic cells. Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.




Western blot analysis of Hela with TDY136 diluted at

1) 1:2,000

2) 1:5,000



Western blot analysis of 1) Rat Testis Tissue, 2) Raw264.7 with TDY136 diluted at 1:2000




1. Input: Hela Cell Lysate

2, IP product: IP dilute 1:200

Western blot analysis: primary antibody: TDY136 1:1,000
Secondary antibody: Goat anti-Mouse IgG(H+L) (S001) 1:10,000



天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.



Immunohistochemical analysis of paraffin-embedded Human Clolon using Histone H3(tri methyl K9) (TDY136) Mouse mAb diluted at 1:500.