

天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

Histone H3(Phospho Thr6) Rabbit Polyclonal Antibody

Catalog TDY228C TDY228F Tel: 010-82908854

Free: 400-0620-621

Quantity 50µL 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC	H, R, M	~15-17KD	IqG

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:1,000 IHC: 1:200-500

Optimal dilutions should be determined by the end user.


Specificity: The Histone H3(Phospho Thr6) Rabbit Polyclonal Antibody can detects endogenous Histone H3 (Phospho Thr6) protein.

Alternative Names: H3 histone antibody, HIST1H3A antibody, Histone cluster 1, H3a antibody

Background: Histone H3 is one of the five main histone proteins involved in the structure of chromatin in eukaryotic cells. Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.



Nocodazole



Western blot analysis of extracts from Hela cells, untreated (-) or treated with Nocodazole (1µg/ml, 24 hr; +), using Phospho-Histone H3 (Thr6)Rabbit pAb (upper) TDY228 1:1000 and Histone H3 (1G1) Mouse mAb (lower) TDY055 1:2000.

Immunohistochemical analysis of paraffin-embedded Human Breast Caricnoma using Histone H3(Phospho Thr6) (TDY228) Rabbit pAb diluted at 1:500.

Applications: WB-Western blot IHC-Immunochemistry IF-Immunofluorescence IP-Immunoprecipitation ChIP-Chormatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig Z-Zebrafish