

Histone H4(Mono Methyl Arg3) Rabbit Polyclonal Antibody

| Catalog | TDY292C | TDY292F | | Tel: 010-82908854 |
|------------------------|---------|--------------------------|------------------|---------------------|
| | | | | Free: 400-0620-621 |
| Quantity | 50µL | 100µL | | Web: www.tdybio.com |
| For research use only. | | | | |
| Applications | | Species Cross-Reactivity | Molecular Weight | Isotype |
| WB, IHC | | Н | 14KD | lgG |

Storage Buffer & Condition: Immunogen Affinity Purified Rabbit IgG, 1mg/ml in PBS pH7.4, containing 0.02% *sodium azide* as Preservative and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:1,000-2,000 IHC: 1:200-500


Optimal dilutions should be determined by the end user.

Specificity: The Histone H4(Mono Methyl Arg3) Rabbit Polyclonal Antibody can detects endogenous Histone H3 (Mono Methyl Arg3) protein.

Alternative Names: Histone 1 H4 antibody, HIST1H4K antibody, MGC24116 antibody, dJ160A22.1 antibody

Background: Histone H4 is one of the 5 main histone proteins involved in the structure of chromatin in eukaryotic cells. H4 is a structural component of the nucleosome, and is subject to covalent modification, including acetylation and methylation, which may alter expression of genes located on DNA associated with its parent histone octamer.





Western blot analysis of extracts from Hela cells, using Histone H4(Mono Methyl Arg3) Rabbit pAb (upper) TDY292 1:2,000.

Immunohistochemical analysis of paraffin-embedded Human Breast Caricnoma using Histone H4 (Mono Methyl Arg3) (TDY292) Rabbit pAb diluted at 1:500.