

天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

Histone H3(Tri Methyl Lys18) Rabbit Polyclonal Antibody

Catalog TDY307C TDY307F Tel: 010-82908854

Free: 400-0620-621

Quantity 50µL 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC	H, M	15-17KD	IgG

Storage Buffer & Condition: Antigen Affinity Purified Rabbit IgG, 1mg/ml in PBS pH7.4, containing 0.02% **sodium azide** as Preservative and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:500-1,000 IHC: 1:200-500

Optimal dilutions should be determined by the end user.


Specificity: The Histone H3(Tri Methyl Lys18) Rabbit Polyclonal Antibody can detects endogenous Histone H3 (Tri Methyl Lys18) protein.

Alternative Names: H3 histone antibody, HIST1H3A antibody, Histone cluster 1, H3a antibody

Background: Histone H3 is one of the five main histone proteins involved in the structure of chromatin in eukaryotic cells. Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.



Western blot analysis of 1) Hela, 2)3T3 with Histone H3 (Tri Methyl Lys18) Rabbit pAb TDY307 diluted at 1:1,000.



Immunohistochemical analysis of paraffin-embedded Human Breast Caricnoma using Histone H3 (Tri Methyl Lys18) (TDY307) Rabbit pAb diluted at 1:500.