

STAT3 Mouse Monoclonal Antibody(1E3)

Catalog TDY477C TDY477F

Tel: 010-80117836

Web: www.tdybio.com

Quantity 50μL 100μL

Entrez-Gene ID#6774 , Swiss-Prot Acc.#P40763

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB,IHC	H,R,M	79,86KD	IgG1

Storage Buffer & Condition: PBS, pH 7.4, containing 0.02% **sodium azide** as Preservative and 50% Glycerol.

Store at **-20°C. Do not aliquot the antibody.**


Recommended dilutions: WB: 1:1,000-2,000 IHC: 1:100-200

Optimal dilutions should be determined by the end user.


Specificity: Antibody can detects endogenous STAT3 protein.

Alternative Names: APRF antibody, HIES antibody, Signal transducer and activator of transcription 3 antibody


Background: Signal transducer and activator of transcription 3, also known as STAT3, is a transcription factor which in humans is encoded by the *STAT3* gene. Stat3 is constitutively activated in a number of human tumors and possesses oncogenic potential and anti-apoptotic activities.



Western blot analysis of 1) Hela, 2)3T3, 3) PC12 with STAT3 Mouse mAb TDY477 diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Human Colon Carcinoma using STAT3 (TDY477) Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Brain Tissue

using STAT3 (TDY477) Mouse mAb diluted at 1:200.

Applications: WB-Western blot IHC-Immunochemistry IF-Immunofluorescence IP-Immunoprecipitation ChIP-Chormatin Immunoprecipitation
Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig Z-Zebrafish